Soil sample boxes can be picked up at your county MSU Extension Service office. It is important that all information asked for on the container, as well as any additional paperwork, be filled out completely and accurately. Crop code 95 should be used for any wildlife seed mixtures (clover, alfalfa, ryegrass, chufa, etc.).

For more information, forms, sample boxes, sampling publications, packaging, and instructions, contact your local county Extension office or the MSU Extension Service Soil Testing Laboratory. The cost is \$6 per soil sample.

Samples may be sent directly to the Soil Testing Laboratory at the address below or may be returned to your local county Extension office. Be sure to include the information sheet and check or money order with the package. Label and tightly secure the shipping container.

MAIL PACKAGES TO MSU Extension Service Soil Testing Laboratory Box 9610 Mississippi State, MS 39762-9610

The MSU Extension Service Soil Testing Laboratory is committed to meeting the needs of its clientele with an accurate and timely report of each soil or plant sample submitted.





EXTENSION

extension.msstate.edu

Copyright 2012 by Mississippi State University. All rights reserved. This publication may be copied and distributed without alteration for nonprofit educational purposes provided that credit is given to the Mississippi State University Extension Service.

By Karl K. Crouse, Ph.D., Associate Extension Professor, Plant and Soil Sciences, and Ben West, Ph.D., former Assistant Extension Professor, Wildlife and Fisheries.

We are an equal opportunity employer, and all qualified applicants will receive consideration for employment without regard to race, color, religion, sex, national origin, disability status, protected veteran status, or any other characteristic protected by law.

Publication 2363

Extension Service of Mississippi State University, cooperating with U.S. Department of Agriculture. Published in furtherance of Acts of Congress, May 8 and June 30, 1914. GARY B. JACKSON, Director

Soil tests for VILDLIFE



IF YOU ARE INTERESTED IN *enhancing*

wildlife ON YOUR PROPERTY **bigger** *deer*, more *quail*, larger *turkeys*—

THEN YOU MUST UNDERSTAND THE

IMPORTANCE OF SOILS. SOIL FERTILITY and

TYPE INFLUENCES MANY ASPECTS OF

WILDLIFE, INCLUDING POPULATION

DISTRIBUTION, ABUNDANCE, and QUALITY.

FOR EXAMPLE, RESEARCH HAS SHOWN THAT

ANTLER SIZE IN WHITE-TAILED DEER IS

DIRECTLY AFFECTED BY SOIL FERTILITY.

The establishment of food plots is one management technique that more and more landowners are using to benefit wildlife. If you are interested in establishing food plots for wildlife, a soil test is necessary for success. A soil test will provide recommendations for lime and fertilizer, customized to your site and for the plants in which you are interested. The MSU Extension Service Soil Testing Laboratory runs a variety of tests on soil samples, including pH, buffer pH (lime requirement), phosphorus, potassium, calcium, magnesium, sodium, and zinc. Based on these tests, your report will recommend levels of nitrogen, phosphorus, and potassium needed for your specific situation. In addition, a soil test will provide you with lime recommendations to balance acidic soils, which are common in Mississippi. However, because some soils within the state are alkaline and do not require lime, a soil test is always recommended to determine the requirements for your site. Lime can improve the physical, chemical, and biological conditions in acidic soils, resulting in greater root proliferation, earlier aboveground plant growth, and

improved nutrient and water uptake. Without proper liming, fertilization can be a wasted effort and expense (see table).

Proper collection of soil samples is extremely important to ensure reliable recommendations. To properly collect a soil sample, take several sub-samples from throughout the field and mix them together to obtain an average for the entire area. The number of sub-samples needed is dependent on the size of the field; as a rule of thumb, take about 10 sub-samples for every acre in the field. Thus, if you planned to establish a 1-acre wildlife food plot, plan on taking 10 sub-samples that you would combine into a single sample for that field. Sub-samples should be collected from the top layer of soil, which is 0-6 inches in depth. Mix the subsamples together to form a uniform sample and discard any plant material that could have been collected.



SOIL ACIDITY	NITROGEN	PHOSPHATE	POTASH	FERTILIZER WASTED
Extremely Acid 4.5 pH	30%	23%	33%	71.34%
Very Strong Acid 5.0 pH	53%	34%	52%	53.67%
Strongly Acid 5.5 pH	77%	48%	77%	32.69%
Medium Acid 6.0 pH	89%	52%	100%	19.67%
Neutral 7.0 pH	100%	100%	100%	0.00%